arXiv:math/0207043v1 [math.DS] 4 Jul 2002

On quasi-invariant transverse measures for the horospherical foliation of a negatively curved manifold

Barbara SCHAPIRA

MAPMO, Université d'Orléans, Rue de Chartres,
BP 6759, 45067 Orléans cedex 2, France
schapira@labomath.univ-orleans.fr

February 1, 2008

Abstract

If M is a compact or convex-cocompact negatively curved manifold, we associate to any Gibbs measure on T^1M a quasi-invariant transverse measure for the horospherical foliation, and prove that this measure is uniquely determined by its Radon-Nikodym cocycle. (This extends the Bowen-Marcus unique ergodicity result for this foliation.) We shall also prove equidistribution properties for the leaves of the foliation w.r.t. these Gibbs measures. We use these results in the study of invaiant meausres for horospherical foliations on regular covers of M.

Primary AMS classification: 37D40, 37C85, 37A20, 22F05

Secondary AMS classification: 37D35, 32Q05, 58Hxx,

Keywords: horospherical foliation, quasi-invariant measures, unique ergodicity, Gibbs measures.

1 Introduction and statement of results

Let $M = \Gamma \backslash \widetilde{M}$ be a complete riemannian manifold with pinched negative curvature, with universal cover $\widetilde{M}$ and fundamental group Γ. Then the unit tangent bundle $T^1\widetilde{M}$ of $\widetilde{M}$ carries the *horospherical foliation* $\widetilde{\mathcal{W}}^{su}$, whose leaves are the unit vectors normal to horospheres and pointing outward. Passing to the quotient leads to a foliation $\mathcal{W}^{su}$ of T^1M which is the strong unstable foliation of the geodesic flow of M. In this paper, we shall construct quasi-invariant measures for this foliation, and study their ergodic properties.

Recall that for a compact manifold M, it was proved by Bowen and Marcus that the foliation $\mathcal{W}^{su}$ is uniquely ergodic: there exists, up to a multiplicative constant, a unique transverse measure for the foliation which is invariant under holonomy. Their proof used symbolic dynamics and showed that the transverse measure is induced by the Bowen-Margulis measure m^0 of T^1M, which is also the measure of maximal entropy of the geodesic flow [7].

In the last decade, it was realized that the measure m^0 could be constructed in a more geometrical way using the Patterson measure on the boundary at infinity of $\widetilde{M}$ [18] [17]. This construction allows to go beyond compact manifolds, and, for instance, Roblin was able to give a purely geometrical proof of the unique ergodicity of $\mathcal{W}^{su}$ for a convex-cocompact manifold, under the assumption that the geodesic flow is topologically mixing [26].

In a similar way, one can associate to a Hölder function $f : T^1M \to \mathbf{R}$ a measure m^f on T^1M, which is the equilibrium state of f when M is compact or convex-cocompact, the Bowen-Margulis measure m^0 corresponding to $f \equiv 0$ see e.g. [16][22][9]. We shall first see that this measure m^f induces a *quasi-invariant transverse measure* for the foliation $\mathcal{W}^{su}$. We use here the definition of a quasi-invariant measure of a foliation as introduced by Connes in [8] in relation with the theory of C^*-algebras associated to a foliation. First, we recall that a *cocycle* for a foliation is a map ρ defined on the set of pairs of points in the same leaf such that $\rho(u, v) + \rho(v, w) = \rho(u, w)$, and that a *transverse measure* $\mu = \{\mu_T\}$ associates to any tranversal T to the foliation a Borel measure μ_T supported on T. Then μ is said to be quasi-invariant if there exists a cocycle ρ such that for each holonomy map ζ between two transversals T and T', $\zeta_*\mu_T$ is absolutely continuous with respect to μ'_T with Radon-Nikodym derivative given by

$$\frac{d\zeta_*\mu_T}{d\mu_{T'}}(\zeta x) = \exp(\rho(x, \zeta x)).$$

Note that if $\mu' = \{\mu'_T\}$ is *equivalent* to $\{\mu_T\}$ (i.e. there exists a Borel map $\psi : T^1M \to \mathbf{R}$ such that, for all transversal T, $d\mu_T(v) = \exp\psi(v)\, d\mu'_T(v)$), then the Radon-Nikodym cocycle ρ' of μ' is *cohomologous* to ρ (there exists a Borel map $R : T^1M \to \mathbf{R}$, such that $\rho(v, w) = \rho'(v, w) + R(v) - R(w)$).

Let us call two Hölder functions f and f' *equivalent* if there exists a Hölder map $\phi : T^1M \to \mathbf{R}$, differentiable in the direction of the geodesic flow, and a constant c such that $f = f' + c + X.\phi$, with $X : TM \to TTM$ the geodesic spray.

Our starting point is the following Proposition: it provides a large family of cocycles which are Radon-Nikodym cocycles of quasi-invariant measures.

Proposition 1 *To each Hölder map* $f : T^1M \to \mathbf{R}$ *is associated an explicit Hölder cocycle* ρ^f *for the foliation* $\mathcal{W}^{su}$*, and a transverse measure* $\mu^f = \{\mu^f_T\}$*, which is quasi-invariant with cocycle* ρ^f*. Moreover, the cohomology class of* ρ^f *and the equivalence class of* μ^f *depend only the equivalence class of* $[f]$*.*

For $f \equiv 0$, the cocycle ρ^0 is trivial, and one recovers a transverse invariant measure μ^0. For compact or convex-cocompact manifolds, one knows that this measure is the unique invariant mesure of the foliation, and thus we are led to the following question: is the transverse measure μ^f the unique quasi-invariant measure with the given cocycle ρ^f?
In the sequel of this work, we will always make the following assumptions:

Assumptions: *The fundamental group* $\Gamma = \pi_1(M)$ *is cocompact or convex-cocompact, and in the second case, the geodesic flow is topologically mixing on its nonwandering set.*

Under these assumptions, our main result is:

Theorem 2 *Let* $[\rho^f]$ *be a cohomology class of cocycles associated to a class* $[f]$ *of Hölder maps. Then for all* $\rho \in [\rho^f]$*, there exists, up to a multiplicative constant, a unique transverse measure* μ *quasi-invariant with Radon-Nikodym cocycle* ρ*, and it is equivalent to* μ^f*.*

When $f \equiv 0$ we recover the unique ergodicity of the foliation. Theorem 2 was already proved by Babillot-Ledrappier [4] for a compact manifold using symbolic dynamics, but we give here a geometrical proof, which allows to prove further results as the equidistribution property of leaves of the foliation (see below Theorem 3).

Theorem 2 can be reinterpreted in terms of the action of the fundamental group Γ of M on the boundary at infinity $\partial\widetilde{M}$ of $\widetilde{M}$. Indeed the lifted foliation $\widetilde{\mathcal{W}}^{su}$ of $T^1\widetilde{M}$ admits a nice set of leaves: the space $\mathcal{H}$ of horospheres, which can be parametrized as $\partial\widetilde{M} \times \mathbf{R}$. An invariant transverse measure for $\mathcal{W}^{su}$ lifts to an invariant transverse measure for $\widetilde{\mathcal{W}}^{su}$ which is Γ-invariant, and thus induces in a canonical way a Γ-invariant measure on $\mathcal{H}$. In the case of quasi-invariant measures, such a correspondance can be made by chosing a global tranversal of $\widetilde{\mathcal{W}}^{su}$, and thus depends on this choice. Our construction is slightly different : we associate to each quasi-invariant transverse measure $\{\mu^f_T\}$ a quasi-invariant measure on $\mathcal{H}$ with the additional property that its Radon-Nikodym cocycle

$$c^f : \Gamma \times \mathcal{H} \mapsto \mathbf{R}$$

is in fact a cocycle for the action of Γ on $\partial\widetilde{M}$: for all $(\xi, s) \in \partial\widetilde{M} \times \mathbf{R} \simeq \mathcal{H}$, and $\gamma \in \Gamma$,

$$c^f(\gamma, (\xi, s)) = c^f(\gamma, \xi).$$

To avoid confusions, we will denote by $[c^f]_{\mathcal{H}}$ and $[c^f]_{\partial\widetilde{M}}$ the respective cohomology classes of c^f as a cocycle for the action of Γ on $\mathcal{H}$ and on $\partial\widetilde{M}$.
A result of Ledrappier, [22], shows that any Hölder cocycle c on $\partial\widetilde{M}$ can be written as $c = c^f$, for a convenient Hölder map $f : T^1M \to \mathbf{R}$. Thus we can reformulate Theorem 2 in the following way:

Theorem 2 bis *Let $[c]_{\mathcal{H}}$ be the cohomology class on $\mathcal{H}$ of a Hölder cocycle c on $\partial\widetilde{M}$. There exists, up to a multiplicative constant, a unique class $[\widehat{\mu}]$ of Γ-quasi-invariant measures on $\mathcal{H}$ with respect to this class $[c]_{\mathcal{H}}$, and they are ergodic.*

This result leads to the question whether there exist cohomology classes of cocycles on $\mathcal{H}$ which do not contain cocycles arising from cocycles on $\partial\widetilde{M}$, and if the Γ-quasi-invariant measures on $\mathcal{H}$ with such a cocycle can be characterized.

In the case of a flow, the property of unique ergodicity implies (and is even equivalent to) the equidistribution of all orbits to the unique invariant measure. In our set-up, we have no flow, so we have to consider more general means on leaves of the foliation. Then we shall see that the above implication is still true.

Theorem 3 *Let $f : T^1M \to \mathbf{R}$ be a Hölder map. There exists on each strong unstable horosphere H^+ a measure $\overline{\mu}^f_{H^+}$, and a distance d_{H^+}, such that for all non wandering vector $u \in T^1M$, the mean $M_{r,u}$ on the ball $B^+(u,r)$ w.r.t. the measure $\overline{\mu}^f_{H^+}$ converges weakly to the equilibrium state m^f associated to f when $r \to \infty$.*

We will also see that the equilibrium state m^f is locally equivalent to the product $\widehat{\mu}^f \times \overline{\mu}^f_{H^+}$.

In fact, we shall deduce this Theorem as a corollary of the more general Theorem 8 of equidistribution of sequences of sets $(E_n)_{n\in\mathbf{N}}$ on leaves which satisfy a certain growth property.

As another application of the preceding results, we shall address the problem of determining the $\bar{\Gamma}$-invariant measures on $\mathcal{H}$, with $\bar{\Gamma} \lhd \Gamma$ a normal subgroup of Γ. Babillot and Ledrappier proved in [4] that when Γ is cocompact, and $\Gamma/\bar{\Gamma}$ is isomorphic to $\mathbf{Z}^d$, each cohomology class $[\alpha]$ of 1-forms on M, which vanishes on loops of $\bar{\Gamma}$, induces a $\bar{\Gamma}$-invariant measure on $\mathcal{H}$, and that these measures are $\bar{\Gamma}$-conservative and ergodic. However, it is still an open question to see whether the measures they constructed are the only $\bar{\Gamma}$-invariant and ergodic measures on $\mathcal{H}$. Here is a partial answer under an additional assumption: for a general normal subgroup $\bar{\Gamma}$ of a cocompact or convex-cocompact group Γ, these measures are the only possible Γ-quasi-invariant, $\bar{\Gamma}$-invariant, and $\bar{\Gamma}$-ergodic measures on $\mathcal{H}$.

Theorem 4 *Let $\bar{\Gamma} \lhd \Gamma$ be a normal subgroup. If $\alpha : TM \to \mathbf{R}$ is a closed 1-form vanishing on the image of $\bar{\Gamma}$ in $H_1(M, \mathbf{R})$, then the measure $\widehat{\mu}^{\alpha}$ associated to the*

Hölder map $\alpha_{|T^1M}$ *is* $\bar{\Gamma}$*-invariant. Conversely, any* $\overline{\Gamma}$*-invariant and* $\overline{\Gamma}$*-ergodic measure* $\widehat{\nu}$ *on* $\mathcal{H}$*, which is quasi-invariant by* Γ*, is necessarily a measure of the form* $\widehat{\nu} = \widehat{\mu}^\alpha$*, with* α *a closed* 1*-form (vanishing on* $\overline{\Gamma}$*).*

Note that in the general case $\overline{\Gamma} \lhd \Gamma$, it is not clear whether the measures $\widehat{\mu}^\alpha$ are $\overline{\Gamma}$-ergodic; see [19], [20] for cases where it is known and other references.

The organization of the text is the following: in a first part, we introduce some notations and all the measures that we will study. In part 3, we prove the unicity results (Theorems 2 and 2 bis). We deduce then of these results an equidistribution result for balls and for more general sets on horospheres (Theorem 8 and Theorem 3). Finally, we prove Theorem 4 in part 5.

It is a pleasure to thank Vadim Kaimanovich for his very careful reading, and many useful suggestions on a preliminary version of this work. Also, I would like to thank my supervisor Martine Babillot for her continuous and valuable guidance during the elaboration of this article, and Jean-Pierre Otal for several discussions.

2 Notations, preliminaries

2.1 Geometry

Let $M = \Gamma\backslash\widetilde{M}$ be a complete riemannian manifold with pinched negative sectional curvature, $\widetilde{M}$ its universal cover, and Γ its fundamental group. Then $\widetilde{M}$ can be compactified into $\overline{M} = \widetilde{M} \cup \partial\widetilde{M}$, where $\partial\widetilde{M}$ is the *boundary at infinity* of $\widetilde{M}$, i.e the set of equivalence classes of geodesic rays which stay at bounded distance one another. The group Γ acts on $\widetilde{M}$ by isometries, and on $\partial\widetilde{M}$ by homeomorphisms. The *limit set* Λ of Γ is the set of accumulation points of Γ in $\partial\widetilde{M}$: $\Lambda = \overline{\Gamma x}\backslash\Gamma x$, for any $x \in \widetilde{M}$. We denote by T^1M (resp. $T^1\widetilde{M}$) the unit tangent bundle of M (resp. $\widetilde{M}$), and by $\pi : T^1M \to M$ (resp. $T^1\widetilde{M} \to \widetilde{M}$) the canonical projection. We will use the distance d on M (and $\widetilde{M}$) induced by the riemannian structure.

The geodesic flow $\Phi = (\Phi^t)_{t\in\mathbf{R}}$ (resp. $\widetilde{\Phi}$) associates to a pair $(t, v) \in \mathbf{R} \times T^1M$ (resp. $\mathbf{R}\times T^1\widetilde{M}$) the tangent vector $\Phi^t v = \dot{c}_v(t)$ at time t to the unique geodesic c_v of M such that $\dot{c}_v(0) = v$.

By a theorem of Eberlein [11], the *nonwandering set* $\Omega \subset T^1M$ of the geodesic flow Φ is the set of vectors v such that any lift $\widetilde{v}$ to $T^1\widetilde{M}$ has both endpoints $c_v(\pm\infty)$ in the limit set Λ.

The group Γ is *cocompact* if $M = \Gamma\backslash\widetilde{M}$ is compact, and then $\Lambda = \partial\widetilde{M}$, and $\Omega = T^1M$. It is *convex-cocompact* when Ω is compact. In this work, we assume that Γ is cocompact or convex-cocompact, so Ω will always be a compact set. We assume also that the geodesic flow is *topologically mixing* on its non-wandering set Ω, which is equivalent to the nonarithmeticity of the length spectrum of M.

It is known to be true in the case of a surface, in the constant curvature case, and in some other cases (see Dal'bo, [10]).
A Hölder *cocycle* on $\partial\widetilde{M}\times\widetilde{M}\times\widetilde{M}$ is a Hölder map $c:\partial\widetilde{M}\times\widetilde{M}\times\widetilde{M}\to\mathbf{R}$ such that for all $\xi\in\partial\widetilde{M}$, and x,y,z in $\widetilde{M}$, $c_\xi(x,y)+c_\xi(y,z)=c_\xi(x,z)$. It is said to be Γ-invariant if it is invariant under the diagonal action of Γ on $\partial\widetilde{M}\times\widetilde{M}\times\widetilde{M}$.
The *Busemann cocycle* is defined on $\partial\widetilde{M}\times\widetilde{M}\times\widetilde{M}$ by:

$$\beta_\xi(x,y)=\lim_{z\to\xi}d(x,z)-d(y,z)=\text{"}d(x,\xi)-d(y,\xi)\text{"}.$$

It is a continuous and Γ-invariant cocycle on $\partial\widetilde{M}\times\widetilde{M}\times\widetilde{M}$. By abuse of notation, if v,w are vectors on $T^1\widetilde{M}$ with base points x and y on $\widetilde{M}$, we will often note $\beta_\xi(v,w)$ in place of $\beta_\xi(x,y)$.
The space of oriented geodesics can be identified to the *double boundary*, i.e. the set of pairs of distinct points $(\eta_1,\eta_2)\in\partial^2\widetilde{M}:=\partial\widetilde{M}\times\partial\widetilde{M}\backslash\{(\xi,\xi),\xi\in\partial\widetilde{M}\}$. The Busemann cocycle allows to give coordinates on $T^1\widetilde{M}$ in terms of this double boundary. More precisely, if a point $o\in\widetilde{M}$ is fixed, the map defined by:

$$v\in T^1\widetilde{M}\to(v^-,v^+,\beta_{v^-}(v,o))\in\partial^2\widetilde{M}\times\mathbf{R}$$

is an homeomorphism.



Figure 1: Coordinates on $T^1\widetilde{M}$

On $\partial^2\widetilde{M}\times\mathbf{R}$, the actions of the geodesic flow and the group Γ commute and can be written in the following way: for any $\gamma\in\Gamma$,

$$\gamma(v^-,v^+,s)=(\gamma v^-,\gamma v^+,s+\beta_{v^-}(o,\gamma^{-1}o))\quad\text{and}\quad\widetilde{\Phi}^t(v^-,v^+,s)=(v^-,v^+,s+t).$$

Thus, the unit tangent bundle T^1M identifies to the quotient $\Gamma\backslash(\partial^2\widetilde{M}\times\mathbf{R})$, and the non-wandering set Ω to $\Gamma\backslash(\Lambda^2\times\mathbf{R})$.
A *horosphere* $H\subset\widetilde{M}$ centered at ξ is a level set of a function $x\to\beta_\xi(x,y)$. These horospheres lift to $T^1\widetilde{M}$: if $u\in T^1\widetilde{M}$, and H is the horosphere centered at u^- and containing the base point $\pi(u)\in\widetilde{M}$ of u, the *strong unstable horosphere of* u, denoted by $H^+(u)$ is the set of vectors $v\in T^1\widetilde{M}$ such that $v^-=u^-$ and $\pi(v)\in H$. It is also the set of vectors of $T^1\widetilde{M}$ with base points in H, orthogonal



Figure 2: Horospheres

to H and pointing outwards. Similarly, we denote by $H^-(u)$ the strong stable horosphere of u. We denote by $\mathcal{H}$ the *space of horospheres.*
The homeomorphism $T^1\widetilde{M} \simeq \partial^2\widetilde{M} \times \mathbf{R}$ allows to identify a horosphere $H^+(u)$ with $\{u^-\} \times \partial\widetilde{M}\backslash\{u^-\} \times \{s(u)\}$, and the space $\mathcal{H}$ of horospheres with $\partial\widetilde{M} \times \mathbf{R}$. The group Γ acts on $\mathcal{H}$ by $\gamma(\xi, s) = (\gamma\xi, s+\beta_\xi(o, \gamma^{-1}o)$. Besides, the horospheres $H^+(u)$ are exactly the strong unstable manifolds of the geodesic flow $\widetilde{\Phi}$:

$$H^+(u) = \widetilde{W}^{su}(u) := \{w \in T^1\widetilde{M},\ \lim_{t\to-\infty} D(\widetilde{\Phi}^t u, \widetilde{\Phi}^t w) = 0\}.$$

Here, D is the distance on $T^1\widetilde{M}$ induced by the Sasaki metric on $T\widetilde{M}$. Similarly, the strong stable horospheres $H^-(u)$ equal the strong stable manifolds $\widetilde{W}^{ss}(u)$. These strong unstable horospheres form a foliation $\widetilde{\mathcal{W}}^{su}$ of $T^1\widetilde{M}$, called the *horospherical foliation*, or strong unstable foliation. A natural family of transversals to $\widetilde{\mathcal{W}}^{su}$ is the family of weak stable manifolds:

$$\widetilde{W}^s(v) = \{w \in T^1M, \exists C = C_w > 0, \forall t \geq 0, D(\Phi^t v, \Phi^t w) \leq C\}.$$

Viewed on $\partial^2\widetilde{M} \times \mathbf{R}$, a transversal $\widetilde{W}^s(v)$ equals $\partial\widetilde{M}\backslash\{v^+\} \times \{v^+\} \times \mathbf{R}$. Recall that, given two (small) transversals T, T' to the foliation included in the same chart of the foliation, a *holonomy map* is a homeomorphism from an open subset of T to T', which preserves the leaves of the foliation. Note that there are natural (global) holonomy maps between two such transversals, given by:

$$\begin{array}{rcl} \widetilde{W}^s(v) & \xrightarrow{\zeta} & \widetilde{W}^s(w) \\ u = (u^-, v^+, s(u)) & \mapsto & \zeta(u) = (u^-, w^+, s(u)). \end{array}$$

The foliation $\widetilde{\mathcal{W}}^{su}$ induces on the quotient $\widetilde{M}/\Gamma$ the foliation $\mathcal{W}^{su}$ whose leaves are the strong unstable manifolds W^{su} for Φ. For the latter foliation, the holonomy maps are not defined globally.

Finally, let us introduce useful families of distances. First, the family $(d_x)_{x\in\widetilde{M}}$ of Gromov distances on the boundary is defined by:

$$\forall(\xi, \eta) \in \partial^2\widetilde{M},\ d_x(\xi, \eta) = \exp\left(-\frac{1}{2}\beta_\xi(x, y) - \frac{1}{2}\beta_\eta(x, y)\right),$$

with y any point on the geodesic (ξ,η). (Rigorously, they are known to be true distances if the curvature of M is less than or equal to -1 [5]; otherwise, one has to take convenient powers of the above quantities to obtain distances.) Using these distances, Hamenstädt [15] defined on each horosphere H^+ a distance d_{H^+} by the formula:

$$\forall (u,v)\in (H^+)^2,\ d_{H^+}(u,v)=\exp\left(\frac{1}{2}\beta_{u^+}(x,u)+\frac{1}{2}\beta_{v^+}(x,v)\right)\, d_x(u^+,v^+)\,,$$

where $x\in \widetilde{M}$ is any point (one easily checks that the above formula does not depend on x). Remark that these distances can also be defined on the strong stable horospheres $H^-(v)$. The following picture shows what they represent geometrically: if $u,v\in H^+$, $d_{H^+}(u,v)=\exp(\pm d(a_u,a_v)/2)$ where a_u and a_v are the respective intersection points of $H^-(u)$ and $H^-(v)$ with the geodesic (u^+,v^+) (the sign depends on the order of a_u and a_v on (u^+,v^+)).



$d_{H^+}(u,v)=\exp(-d(a_u,a_v))$ $\qquad$ $d_{H^+}(u,v)=\exp(d(a_u,a_v))$

Figure 3: Horospherical distance

Two important properties of this family are that they are invariant by isometries:

$$\forall \gamma\in\Gamma,\ d_{\gamma H^+}(\gamma u,\gamma v)=d_{H^+}(u,v),$$

and that they “explode” when pushed by the flow:

$$\Phi^t B^+(u,r)=B^+(\Phi^t u, re^t)$$

2.2 Cocycles and measures associated to a Hölder map

A *Hölder potential* is a map $f: T^1M\to \mathbf{R}$ which is Hölder w.r.t. the Sasaki metric D on T^1M. For simplicity, we shall consider for the moment only symmetric potentials: $f(v)=f(-v)$ (see section 5.1 for the general case), and we denote also by f its Γ-invariant lift on $T^1\widetilde{M}$. If x and y are points in $\widetilde{M}$, we denote by $\int_x^y f$ the integral of f on the unique (oriented) geodesic from x to y, viewed on $T^1\widetilde{M}$. If $x,y\in\widetilde{M}$ and $\xi\in\partial\widetilde{M}$, we define the following cocycle:

$$\rho^f_\xi(x,y)=\lim_{t\to+\infty}\int_x^{c(t)} f-\int_y^{c(t)} f\ =\text{''}\int_x^{\xi} f-\int_y^{\xi} f\text{''},$$

where $(c(t))_{t\geq 0}$ is the geodesic ray $[x,\xi)$. This function is well-defined, since when f is Hölder, the difference between the two integrals converges. Moreover, it is Γ-invariant. It is a generalisation of the Busemann cocycle, since when $f\equiv 1$, $\rho^1_\xi(x,y)=\beta_\xi(x,y)$. Moreover, it induces a cocycle on the strong unstable foliation $\widetilde{\mathcal{W}}^{su}$: if v and w are two vectors on the same strong unstable horosphere $H^+(v)$, this cocycle is defined by: $\rho^f(v,w)=\rho^f_{v^-}(v,w)$.
Note that when $f\equiv 1$, $\rho^f\equiv 0$. By Γ-invariance, one may also define this cocycle for the strong unstable foliation $\mathcal{W}^{su}$ on T^1M; indeed, if v and w are on the same leaf of T^1M, the quantity $\rho^f(\widetilde{v},\widetilde{w})$ is Γ-invariant, so it does not depend on the lifts $\widetilde{v},\widetilde{w}$ chosen on the same leaf of $\widetilde{\mathcal{W}}^{su}$. We can also define it directly on T^1M by

$$\rho^f(v,w)=\lim_{t\to-\infty}\int_v^{\Phi_t v} f-\int_w^{\Phi_t w} f.$$

Finally, if we choose an origin $o\in\widetilde{M}$, we can associate to f the following Hölder cocycle on $\partial\widetilde{M}$:

$$c^f_o(\gamma,\xi)=-\rho^f_\xi(o,\gamma^{-1}o).$$

Lemma 2.1 *The cohomology class $[c^f_o]_{\partial\widetilde{M}}$ depends on $f\in[f]$, but on $\mathcal{H}$, the class $[c^f_o]_{\mathcal{H}}$ depends only on $[f]$, and neither on o, nor on $f\in[f]$.*

The proof is an easy computation. We will see after the following proposition that there is another interesting cocycle on $\partial\widetilde{M}$ associated to f.
To each Hölder potential f on T^1M is associated a unique equilibrium state m^f, that is a probability measure on $\Omega\subset T^1M$ satisfying the variational principle (see [22], [9] for the construction, and [18], [16] to see that it is an equilibrium state). This measure is induced on the quotient $T^1M=\Gamma\setminus T^1\widetilde{M}$ by a measure $\widetilde{m}^f$ on $T^1\widetilde{M}$. We will recall the two main steps of its construction.

Proposition 5 (Ledrappier, [22]) *There exists a (unique) probability measure ν^f_o on $\Lambda\subset\partial\widetilde{M}$, which has no atoms, and is ergodic and Γ-quasi-invariant with respect to the following cocycle:*

$$(\gamma,\xi)\to\delta^f\beta_\xi(o,\gamma^{-1}o)-\rho^f_\xi(o,\gamma^{-1}o).$$

The constant δ^f is the topological pressure of f. Moreover the equivalence class $[\nu^f_o]$ of ν^f_o and the cohomology class on $\partial\widetilde{M}$ of its Radon-Nikodym cocycle $\delta^f\beta_\xi(o,\gamma^{-1}o)-\rho^f_\xi(o,\gamma^{-1}o)$ depend only on the equivalence class $[f]$, but neither on o nor on f. More precisely, if o' is another point, we have:

$$\forall\xi\in\Lambda,\quad \frac{d\nu^f_o}{d\nu^f_{o'}}(\xi)\;=\;\exp(\delta^f\beta_\xi(o',o)-\rho^f_\xi(o',o))\,.$$

And if $f=f'+c+X.\phi$, up to a multiplicative constant, we have:

$$\forall\xi\in\Lambda,\quad \frac{d\nu^f_o}{d\nu^{f'}_o}(\xi)=\exp(\phi(v_{o,\xi}))\,,$$

with $v_{o,\xi}$ the vector based at o pointing to ξ.

For simplicity, we will write $\beta^f_\xi(o,\gamma^{-1}o) = \delta^f \beta_\xi(o,\gamma^{-1}o) - \rho^f_\xi(o,\gamma^{-1}o)$. It is another cocycle on $\Gamma \times \partial\widetilde{M}$ associated to f, the Radon-Nikodym cocycle of ν^f_o. Notice also that $\beta^f = \rho^{\delta^f - f}$, so that $[\beta^f]_{\mathcal{H}} = [c^f_o]_{\mathcal{H}}$.
Conversely, Ledrappier characterized the Hölder cocycles which are the Radon Nikodym cocycles associated to a quasi-invariant measure: more precisely, he constructed a bijection between the equivalence classes of Hölder potentials on T^1M, the equivalence classes of quasi-invariant measures on $\Lambda \subset \partial\widetilde{M}$ with a Hölder cocycle, and the cohomology classes on $\partial\widetilde{M}$ of normalized Hölder cocycles.

Recall that we assume for the moment that f is symmetric. The second step is:

Proposition 6 *The Radon measure $\widetilde{m}^f$ on $\Lambda^2 \times \mathbf{R}$ defined below is invariant by Γ and by the geodesic flow:*

$$d\widetilde{m}^f(v) = \exp\left(\beta^f_{v^+}(o,v) + \beta^f_{v^-}(o,v)\right)\, d\nu^f_o(v^-)\, d\nu^f_o(v^+)\, ds.$$

So it induces on the quotient a finite measure on T^1M (normalized to be a probability) with support in the non-wandering set Ω: the equilibrium state m^f associated to f (see [18], [16]). Moreover, this measure is mixing w.r.t the geodesic flow Φ, [2]. Finally, it depends only on the equivalence class $[f]$ of f.

The quasi-product structure of $\widetilde{m}^f$ suggests to decompose it into two families of measures. First, the family $(\mu^f_{H^+})_{H^+\in\mathcal{H}}$ defined by:

$$d\mu^f_{H^+(u)}(v) = \exp(\beta^f_{v^+}(o,v))\, d\nu^f_o(v^+)\,,$$

is a Γ-invariant family of measures on the leaves of the foliation, in the sense that $\gamma_*\mu^f_{H^+} = \mu^f_{\gamma H^+}$.
Recall that a quasi-invariant transverse measure for a foliation is completely determined by its restriction to a family of transversals passing at all points. Then, we have:

Proposition 1' *The transverse measure to the foliation $\widetilde{\mathcal{W}}^{su}$ defined on each transversal $T = \widetilde{W}^s(w)$ by:*

$$\forall v \in T, \quad d\mu^f_T(v) = \exp(\beta^f_{v^-}(o,v))\, d\nu^f_o(v^-)\, ds$$

is invariant by Γ in the sense that $\gamma_*\mu^f_T = \mu^f_{\gamma T}$. So it induces on T^1M a transverse measure to $\mathcal{W}^{su}$ (still denoted by $\{\mu^f_T\}$), which is quasi-invariant by holonomy, with respect to the cocycle ρ^f.

The proof of Proposition 1' is straightforward.

Remark: The above transverse measure $\{\mu^f_T\}$ is constructed using the equilibrium state m^f of f. Thus we can make the analogy between our geometric situation and the symbolic one. In the case of the full shift $\Sigma = \{1,...,k\}^{\mathbf{Z}}$ endowed with the shift σ, the unique equilibrium state m^f coincides with the unique *Gibbs measure* associated to f, that is the measure satisfying $\zeta_* m^f = \exp(\rho^f)\, m^f$ (see [6], [21]), with ζ a map from a cylinder to another. Since a cylinder can be considered as a transversal to the strong stable manifolds of σ, the map ζ can be viewed as a holonomy map. Thus Theorem 2 gives a similar result in the case of a manifold: the unique quasi-invariant transverse measure with respect to ρ^f is "induced" by the unique equilibrium state m^f, which can therefore be called a Gibbs measure.

In our geometrical frame, recall that there exists a nice set of leaves of the foliation: the set $\mathcal{H} \simeq \partial\widetilde{M} \times \mathbf{R}$ of horospheres. Better than working with the family of measures $\{\mu_T\}$ defined above, we would like to obtain a unique measure on $\mathcal{H}$. To do this, we need a holonomy invariant family of measures, or equivalently measures μ_T on transversals $T = \widetilde{W}^s(w)$ which do not depend on T, i.e. on v^+. The only term in the expression of μ_T which is not invariant by holonomy is $\exp(-\rho^f_{v^-}(o,v))$ (when f is not a constant). So we divide the measures μ_T by the density $\exp(-\rho^f_{v^-}(o,v))$, and we multiply the measures $\overline{\mu}^f_{H^+u}$ by the same density. Then we obtain new families of measures:

$$d\overline{\mu}^f_T(v) = ds\, d\nu^f_o(v^-)\, \exp\left(\delta^f \beta_{v^-}(o,v)\right) \qquad \text{and}$$

$$d\overline{\mu}^f_{H^+u}(v) = d\nu^f_o(v^+)\, \exp\left(\beta^f_{v^+}(o,v) - \rho^f_{v^-}(o,v)\right).$$

These families are now quasi-invariant by Γ. More precisely, the measures on leaves satisfy:

$$\forall v \in \Lambda^2 \times \mathbf{R},\ \forall \gamma \in \Gamma, \quad \frac{d\gamma_*^{-1}\overline{\mu}^f_{H^+(\gamma v)}}{d\overline{\mu}^f_{H^+v}}(v) = \exp(-c^f_o(\gamma, v^-)), \tag{1}$$

Pushing by the flow leads to the property:

$$\forall v \in \Lambda^2 \times \mathbf{R},\ \forall t \in \mathbf{R}, \quad \frac{d\Phi_*^{-t}\bar{\mu}^f_{H^+(\Phi^t v)}}{d\bar{\mu}^f_{H^+v}}(v) = \exp(t\delta^f). \tag{2}$$

The family $\overline{\mu}^f_T$ is also quasi-invariant by Γ, with the opposite cocycle: $c^f_o(\gamma, v^-) = \rho^f_{v^-}(\gamma^{-1}o, o)$. But it is now invariant by holonomy, so it induces a measure $\widehat{\mu}^f_o$ on the quotient on the space $\mathcal{H}$ of leaves. More precisely, we have:

Lemma 2.2 *The measure $d\widehat{\mu}^f_o(\xi, s) = \exp(-\delta^f s)\, ds\, d\nu^f_o(\xi)$ on $\mathcal{H}$ is supported on $\Lambda \times \mathbf{R}$, and Γ-quasi-invariant with respect to the cocycle c^f_o.*

Remark that in the case of a constant potential, we did not change anything, these two families of measures are the same as in the beginning, and in particular, the family (μ^f_T) is at the same time Γ-invariant and invariant under the holonomy pseudogroup.

As the topological pressure satisfies $\delta^{f+c} = \delta^f + c$, we can assume that $\delta^f > 0$. It is simply a normalisation convention which allows us to assume that, when pushed by the flow Φ^t, with $t > 0$, the family of measures $\bar{\mu}^f_{H^+}$ is expanded by a factor $\exp(t\delta^f)$.

Let us now summarize the main properties of the above measures in a table.

Measures and their support	Γ-Quasi-Invariance	Q-I by holonomy
ν^f_o on $\Lambda \subset \partial\widetilde{M}$	$\exp(\beta^f)$	—
$\mu^f_{H^+}$ on leaves H^+ of $\mathcal{W}^{su}$	Invariant	—
$\bar{\mu}^f_{H^+}$ on leaves H^+ of $\widetilde{\mathcal{W}}^{su}$	$\exp(-c^f_o)$	—
μ^f_T on transversals to $\mathcal{W}^{su}$	Invariant	$\exp(\rho^f)$
$\bar{\mu}^f_T$ on transversals to $\widetilde{\mathcal{W}}^{su}$	$\exp(c^f_o)$	Invariant
$\widehat{\mu}^f_o$ on $\Lambda \times \mathbf{R} \subset \mathcal{H}$	$\exp(c^f_o)$	—

3 Horospherical means, Proofs of Theorems 2 and 2 bis

The aim of this section is the proof of the unicity results of a quasi-invariant measure (Theorems 2 and 2 bis). We use an intermediate result of equidistribution of horospherical means pushed by the flow Φ^t when $t \to \infty$ [4, thm 4], that we recall below, and some properties of continuity of the measure of balls, proved in section 3.1. A formula (section 3.2) allows in section 3.3 to prove theorem 2 bis. Theorem 2 is obtained as a corollary of the preceding.

Finally, we give in section 3.4 an alternative method to prove Theorems 2 bis and 3 in a particular case. In fact, under the aditional assumption that the boundary of all horospherical balls has measure zero, we are able to prove directly a uniform equidistribution property (Theorem 3 bis), and to deduce easily Theorem 2 bis. (In the general case, recall that we will deduce in section 4 Theorem 3 from Theorem 2 bis.)

Given a potential f, the mean $M_{r,u}(\psi)$ of a map $\psi : T^1M \to \mathbf{R}$ on $B^+(u,r)$ is defined by:

$$M_{r,u}(\psi) \;=\; \frac{1}{\bar{\mu}^f_{H^+u}(B^+(u,r))} \int_{B^+(u,r)} \widetilde{\psi}\, d\bar{\mu}^f_{H^+u}\,,$$

with $\widetilde{\psi}$ the Γ-invariant lift of ψ on $T^1\widetilde{M}$. Notice that these means cannot be defined directly on T^1M, since the measure $\bar{\mu}^f_{H^+u}$ is not Γ-invariant. However, they are Γ-invariant, so we shall consider them as probability measures on T^1M. For simplicity, we denote by $M^t_{r,u}(\psi) := \Phi^t_* M_{r,u}(\psi)$ the mean pushed by the flow. The geometrical property of the horospherical distances then implies the simple, but fundamental property that the mean of $\psi \circ \Phi^t$ on a ball $B^+(u,r)$ equals the mean of ψ on the bigger ball $B^+(\Phi^t u, re^t)$:

$$\forall u \in T^1\widetilde{M},\ \forall r > 0,\ \forall t \in \mathbf{R}, \quad M^t_{r,u}(\psi) = M_{re^t,\Phi^t u}(\psi). \tag{3}$$

Recall also that, from a general result in [4], we have:

Theorem 7 (Babillot, [4]) *Assume that Γ is cocompact or convex-cocompact, and the geodesic flow is topologically mixing on its nonwandering set Ω. For all $u \in \Lambda^2 \times \mathbf{R}$, for all $r > 0$, the mean $M^t_{r,u}$ converges weakly to m^f when $t \to +\infty$.*

This result is a consequence of the mixing of the measure m^f, which is established when the geodesic flow is topologically mixing.
Notice that, after fixing a positive and continuous map $\psi : T^1M \to \mathbf{R}$, we will often consider these means $M^t_{r,u}(\psi)$ as *functions of the variable* $u \in \Lambda^2 \times \mathbf{R}$. By Γ-invariance, we consider them either as functions on Ω or $\Lambda^2 \times \mathbf{R}$.

3.1 Averaging on horospherical balls

In this first section, we shall study the map

$$(u, r) \mapsto \int_{B^+(u,r)} \psi \, d\bar{\mu}^f_{H^+}.$$

It is not continuous, but we shall prove that it has a regularity property. The main result is Lemma 3.3. This paragraph is inspired from the sequence of "technical lemmas" of Roblin, [26], paragraph 1H.
If $u \in T^1\widetilde{M}$ and $v \in H^-(u)$, let us introduce the map $P_{u,v} : H^+(u) \to H^+(v)$ (see Figure 4 below), which sends $w = (u^-, w^+, s(u)) \in H^+(u)$ to the intersection w' of the geodesic (v^-, w^+) with $H^+(v)$. In other words, $w' = (v^-, w^+, s(v)) \in H^+(v)$ (when it makes sense, i.e. when $w^+ \neq u^-, v^-$).



Figure 4: The map $P_{u,v} : w \to w'$

In the following, we prove some lemmas which express mainly three important properties; first, these maps $P_{u,v}$ are uniformly close to the identity when v is close to u (Lemma 3.1). Second, all these measures $\overline{\mu}^f_{H^+}$ have a continuous (and positive) Radon-Nikodym derivative with respect to the measure ν^f_o on $\partial\widetilde{M}$, which allows to compare the measures $\overline{\mu}^f_{H^+(v)}$ and $P_{u,v*}\overline{\mu}^f_{H^+(u)}$: they are equivalent with continuous Radon-Nikodym derivative (Lemma 3.2). Third, the flow contracts the stable manifolds. All these properties are proved in some preliminary lemmas, and collected in Lemma 3.3, which will allow to prove Theorem 2 bis.

To make the discussion more precise, we have to introduce neighborhoods, or "cells", in which we will let v vary: if $u \in T^1\widetilde{M}$, and r_1, r_2, r_3 are three positive numbers, we note

$$C(u, r_1, r_2, r_3) = \cup_{|s|<r_3} \Phi^s \cup_{v_1 \in B^-(u,r_1)} P_{u,v_1}(B^+(u, r_2)).$$



Figure 5: Cells

If ψ is a positive Borel function on $T^1\widetilde{M}$, we set:
$\psi_\varepsilon(w) = \sup\{\psi(v),\ v \in \cup_{|s|<\varepsilon} \Phi^s B^-(w, \varepsilon)\}$, and
$\psi_{-\varepsilon}(w) = \inf\{\psi(v),\ v \in \cup_{|s|<\varepsilon} \Phi^s B^-(w, \varepsilon)\}$. Then, if $A \subset T^1\widetilde{M}$ is a Borel set, we have $[\mathbf{1}_A]_{\pm\varepsilon} = \mathbf{1}_{A_{\pm\varepsilon}}$, and for all $\varepsilon > 0$, and $t \geq 0$, $[\Phi^{-t}A]_\varepsilon \subset \Phi^{-t}(A_\varepsilon)$, and $[\Phi^{-t}A]_{-\varepsilon} \supset \Phi^{-t}(A_{-\varepsilon})$. This means simply that if $t \geq 0$, the flow Φ^t contracts the stable manifolds, and expands the unstable manifolds.

In the first lemma, we compare balls $B^+(u, r)$ and $B^+(v, r)$ when $v \in H^-(u)$.

Lemma 3.1 (Roblin, [26]) *Let $K \subset T^1\widetilde{M}$ be a compact, and $\varepsilon > 0$. There exists $\delta > 0$ such that for all $u \in K$, if $v \in B^-(u, \delta)$ and $w \in B^+(u, 3)$, then $P_{u,v}w \in \Phi^s B^-(w, \varepsilon)$, with $|s| < \varepsilon$, and if $r \in [1, 2]$, $B^+(v, re^{-\varepsilon}) \subset P_{u,v}(B^+(u, r)) \subset B^+(v, re^{\varepsilon})$.*

In other words, the map $P_{u,v} : B^+(u, 3) \to H^+(v)$ restricted to the ball $B^+(u, 3)$ in $H^+(u)$, is uniformly closed to the identity of $B^+(u, 3)$ (up to $\exp(\pm\varepsilon)$) for the topology of uniform convergence on compact sets.

Proof: The assumptions of the Lemma mean that if δ is small enough, the 4-uple (u^-, u^+, v^-, w^+) varies in a compact set of $\partial^2\widetilde{M} \times \partial^2\widetilde{M}$. One easily checks that the quantity s such that $P_{u,v}w \in \Phi^s H^-(w)$ is the cross ratio $B(v^-, u^-, u^+, w^+) =$ "$d(v^-, u^+) + d(u^-, w^+) - d(v^-, w^+) - d(u^-, u^+)$" of the four points. Moreover, the cross ratio is a continuous map, [23], which vanishes if $u^- = v^-$ (or $u^+ = w^+$). Thus if δ is small enough, v^- is uniformly closed to u^-, which gives $P_{u,v}w \in \Phi^s H^-(w)$. Moreover, an easy computation shows that $\dfrac{d_{H^+}(P_{u,v}w, v)}{d_{H^+}(u, v)} = e^{s/2}$, which concludes the proof. $\square$



Figure 6: Cross ratio of (u^-, v^-, u^+, w^+)

The following lemma is a corollary of the above and of continuity of densities of measures $\overline{\mu}^f_{H^+u}$ on horospheres with respect to the measure ν^f_o on the boundary.

Lemma 3.2 *Let $K \subset T^1M$ be a compact, and $\varepsilon > 0$. There exists $\delta > 0$, such that for all $u \in K$, and $v \in B^-(u, \delta)$, for all $w \in B^+(u, 3)$ and $w' = P_{u,v}w$, the following quantity is uniformly close to* 1 *(up to* $\exp(\pm\varepsilon)$*):*

$$\frac{d\overline{\mu}^f_{H^+v}}{d(P_{u,v} * \overline{\mu}^f_{H^+u})}(w') = \exp\left(\beta^f_{w^+}(w, w') - \rho^f_{v^-}(o, w') + \rho^f_{u^-}(o, w)\right).$$

Proof: Recall first that $\beta^f_{w^+}(w, w') - \rho^f_{v^-}(o, w') + \rho^f_{u^-}(o, w) = \delta^f \beta_{w^+}(w, w') - \rho^f_{w^+}(w, w') - \rho^f_{v^-}(o, w') + \rho^f_{u^-}(o, w)$. We have $|\beta_{w^+}(w, w')| \leq \varepsilon$, after 3.1. We have also $|\rho^f_{w^+}(w, w')| \leq C(f)\, D(w, w')^{\alpha(f)}$, because f is Hölder, which gives the desired upper bound by 3.1. Finally, the last term can be written as $\exp(A(P_{u,v}w) - A(w))$, with $A(w) = \rho^f_{w^-}(o, w)$, and the result follows from the continuity of A on $T^1\widetilde{M}$. Indeed, set $w^t = \Phi^{-t}w$, $y \in W^{su}(u)$ the vector of $W^{su}(u)$ tangent to the geodesic $(u^-, o]$, and $y^t = \Phi^{-t}y$. As f is Hölder, and the curvature is bounded above by a negative constant, there exists a $T > 0$, uniform on K, such that if $t \geq T$, $A(w)$ is ε-closed to $\int_o^{y^t} f - \int_w^{w^t} f$. But for all fixed $t > 0$, this quantity is continuous, hence uniformly continuous in $w \in K$. So A is continuous, and the result follows. $\square$

We eventually arrive at the last lemma, which summarizes the main properties of the measure $\overline{\mu}^f_{H^+(u)}$ on balls.

Lemma 3.3 *Let ψ be a positive Borel function on $T^1\widetilde{M}$, $\varepsilon > 0$ (small), and $K \subset T^1M$ a compact set. There exists $r_1, r_2, r_3 > 0$ small enough, so that: for all $u \in K$, for all $v \in C(u, r_1, r_2, r_3)$, $r \in [1, 2]$, we have:*

$$e^{-\varepsilon} \int_{B^+(u, re^{-\varepsilon})} \psi_{-\varepsilon}\, d\overline{\mu}^f_{H^+(u)} \leq \int_{B^+(v,r)} \psi\, d\overline{\mu}^f_{H^+(v)} \leq e^{\varepsilon} \int_{B^+(u, re^{\varepsilon})} \psi_{\varepsilon}\, d\overline{\mu}^f_{H^+(u)}.$$

Sketch of the proof: The proof follows closely that of Roblin, [26], and goes as follows. Let $\varepsilon > 0$, small enough so that $re^{3\varepsilon} \leq 2e^{3\varepsilon} \leq 3$. Fix $u \in K$, and $v \in C(u, r_1, r_2, r_3)$. By definition of cells, we can write $v = \Phi^s v_2$, with $|s| < r_3$, $v_2 \in P_{u,v_1} B^+(u, r_2) \subset H^+(v_1)$ and $v_1 \in B^-(u, r_1)$. First, we push the ball $B^+(v, r)$ by Φ^{-s} to the ball $B^+(v_2, re^{-s})$, with s very small. By relations (2) and (3), the integrals of ψ on these balls are almost the same. Second, we move the ball $B^+(v_2, re^{-s})$ by the map P^{-1}_{u,v_1}. Lemma 3.1 says that the image is uniformly close to the ball $B^+(u, re^{-\varepsilon})$ and Lemma 3.2 says that the derivative of the measure $\mu^f_{H^+}$ is uniformly close to 1. Third, the image of ψ by these operations is close to $\psi_{-\varepsilon}$. The right inequality is proved in the same way. $\square$

3.2 An autoadjonction property

It is obvious that for a flow $(\Phi^t)_{t\in\mathbf{R}}$ on a compact space X, the equality

$$\int_X \frac{1}{T} \int_0^T \psi \circ \Phi^t \, dt d\nu = \int_X \psi \, d\nu$$

for all $T > 0$ and any Φ-invariant measure ν allows to prove unique ergodicity if we know that all Birkhoff averages converge to a constant.
In our situation, the following equality allows to do the same trick, replacing Birkhoff averages by horospherical means: we will use it to prove unique ergodicity, using a weaker property than equidistribution of horospherical means.

Lemma 3.4 *Let $\widehat{\nu}$ be any fixed quasi-invariant measure on $\mathcal{H}$ with the cocycle c^f_o, and M the measure on $\partial^2\widetilde{M} \times \mathbf{R}$ defined by $dM(v) = d\widehat{\nu}(H^+)\, d\overline{\mu}^f_{H^+}(v)$. If ψ a Γ-invariant positive measurable map on $T^1\widetilde{M}$, and D a measurable fundamental domain for the action of Γ on $T^1\widetilde{M}$, then:*

$$\int_D dM(u)\, M_{r,u}(\psi) = \int_D dM(v)\psi(v) \int_{B^+(v,r)} \frac{d\overline{\mu}^f_{H^+(v)}(u)}{\overline{\mu}^f_{H^+(v)}(B^+(u,r))} \, .$$

Before we proceed to the proof, let us make a remark: if we knew that uniform equidistribution of horospherical means when $r \to \infty$ holds, then, as in the case of a flow, the above formula would allow to deduce the unique ergodicity result from the equidistribution property. In fact, we will use this approach in a particular case in section 3.4.

Proof: By definition, $dM = d\widehat{\nu}\; d\overline{\mu}^f_{H^+}$, and the left term can be rewritten:

$$\int_{\mathcal{H}} d\widehat{\nu}(H^+(u)) \int_{H^+(u)\times H^+(u)} d\overline{\mu}^f_{H^+}(u)\, d\overline{\mu}^f_{H^+}(v) \frac{\mathbf{1}_D(u)\, \mathbf{1}_{B^+(u,r)}(v)\, \psi(v)}{\overline{\mu}^f_{H^+(u)}(B^+(u,r))}.$$

The proof is then based on the fact that ψ is Γ-invariant, $\overline{\mu}^f_{H^+}$ and $\widehat{\nu}$ are Γ-quasi-invariant (resp. with cocycles $-c^f_o$ and $+c^f_o$) and on two observations: first

$\mathbf{1}_{B^+(u,r)}(v)$ is symmetric in u and v; second, $B^+(u,r) = \sqcup_{\gamma\in\Gamma} B^+(u,r) \cap \gamma D$. If we set $v = \gamma v'$ and $u = \gamma u'$, we have: $\mathbf{1}_{B^+(u,r)}(v) = \sum_{\gamma\in\Gamma} \mathbf{1}_{B^+(u',r)}(v')\mathbf{1}_D(v')$. If we decompose the above integral into a sum over $\gamma \in \Gamma$, we find that it equals:

$$\sum_{\gamma\in\Gamma} \int_{\mathcal{H}} d\widehat{\nu}(\gamma H^+(u')) \int_{\gamma H^+(u')\times\gamma H^+(u')} d\overline{\mu}^f_{\gamma H^+(u')}(\gamma u')\, d\overline{\mu}^f_{\gamma H^+(u')}(\gamma v') \;\times$$
$$\times\; \frac{\mathbf{1}_{\gamma^{-1}D}(u')\mathbf{1}_{B^+(u',r)}(v')\mathbf{1}_D(v')\psi((\gamma v')}{\overline{\mu}^f_{\gamma H^+(u')}(\gamma B^+(u',r))}$$

We then use the various (quasi-)invariance relations by Γ; we remark also that $\sum_\gamma \mathbf{1}_{\gamma^{-1}D}(u') = 1$, and $H^+(u) = H^+(v)$, and we get:

$$\int_{\mathcal{H}} d\widehat{\nu}(H^+(v)) \int_{H^+(v)\times H^+(v)} d\overline{\mu}^f_{H^+}(v)\, d\overline{\mu}^f_{H^+}(u)\, \frac{\mathbf{1}_D(v)\mathbf{1}_{B^+(v,r)}(u)\psi(v)}{\overline{\mu}^f_{H^+(v)}(B^+(u,r))}.$$

This expression is the same than the first one, with the roles of u and v exchanged. $\square$

3.3 Proof of Theorems 2 and 2 bis

We prove first Theorem 2 bis.

Proof of Theorem 2 bis: Let $[c]_{\mathcal{H}}$ be a cohomology class of Hölder cocycles for the action of Γ on $\mathcal{H}$. By assumption, this class contains a cocycle on $\partial\widetilde{M}$, so we can assume that c is this cocycle. In [22], Ledrappier showed that every Hölder cocycle on $\partial\widetilde{M}$ is of the form $c = \lambda\beta^F$, with F a Hölder potential and $\lambda \in \mathbf{R}$, whence $c = \lambda\rho^{\delta^F - F} = \rho^{\lambda\delta^F - \lambda F}$. Finally, c can be written $c = c^f_o$, with f a Hölder potential.
Let $\widehat{\nu}$ be a nonzero Γ-quasi-invariant measure with the cocycle c^f_o, and M the measure on $\partial^2\widetilde{M} \times \mathbf{R}$ defined by: $dM(v) = d\widehat{\nu}(H^+(v))\, d\overline{\mu}^f_{H^+}(v)$. Let ψ be a positive continuous map on T^1M, and $\widetilde{\psi}$ its lift on $T^1\widetilde{M}$. Lemma 3.4 gives:

$$\int_D dM(u) M^0_{r,u}(\psi) = \int_D dM(v)\widetilde{\psi}(v) \int_{B^+(v,r)} \frac{d\overline{\mu}^f_{H^+(v)}(u)}{\overline{\mu}^f_{H^+(v)}(B^+(u,r))}\ .$$

Since Ω is a compact set, we can choose a relatively compact fundamental domain D for the action of Γ on $\Lambda^2 \times \mathbf{R}$, with $M(\partial D) = \widetilde{m}^f(\partial D) = 0$.
We will prove that for all $r > 0$, the right term is smaller than cst.$\int \psi\, dM$ (inequality (5)), and that there exists $r > 0$, such that the left term is greater than cst$'$.$\int \psi\, dm^f$, so that $m^f \ll M$ (inequality (6)). The constants do not depend on ψ, so that these two inequalities imply $m^f \ll M$.

First, since Γ is cocompact or convex-cocompact, the following Vitali property is satisfied (see [26], Proposition 6.3): there exists a uniform integer N on $C = \Lambda^2 \times \mathbf{R}$, such that for all $u \in C$, and all $r > 0$,

$$B^+(u,r) \cap \overline{C} \subset \cup_{i=1}^N B^+(u_i, \frac{r}{2}). \tag{4}$$

The integral on the ball $B^+(u,r)$ is then smaller than the sum of N integrals on balls $B^+(u_i, \frac{r}{2})$. Moreover, triangular inequality gives $B^+(v,r) \cap \bar{C} \supset B^+(u_i, \frac{r}{2}) \cap \overline{C}$ if $v \in B^+(u_i, \frac{r}{2})$. Hence $\overline{\mu}^f_{H^+(u)}(B^+(v,r)) \geq \overline{\mu}^f_{H^+(u)}(B^+(u_i, \frac{r}{2}))$, and finally,

$$\forall r > 0,\ \forall u \in \Lambda^2 \times \mathbf{R}, \quad \int_{B^+(u,r)} \frac{d\overline{\mu}^f_{H^+(u)}(v)}{\overline{\mu}^f_{H^+(u)}(B^+(v,r))} \leq N.$$

We deduce that:

$$\forall r > 0, \quad \int_D dM(v)\widetilde{\psi}(v) \int_{B^+(v,r)} \frac{d\overline{\mu}^f_{H^+(v)}(u)}{\overline{\mu}^f_{H^+(v)}(B^+(u,r))} \leq N \int_{T^1M} \psi \, dM. \tag{5}$$

We shall now prove the following inequality:

$$\exists r > 0, \exists C > 0, \int_D dM(u) M^0_{r,u}(\psi) \geq C \int_{T^1M} \psi \, dm^f. \tag{6}$$

Let $0 < \varepsilon < \log 2$ be small enough so that $\int_{T^1M} \psi_{-\varepsilon} dm^f \geq \frac{1}{2} \int_{T^1M} \psi \, dm^f$. We can cover D by a finite number of cells $C(u_i)$, $1 \leq i \leq k$, given by Lemma 3.3 for the above ε. If $v \in C(u_i)$, this Lemma gives for all $t \geq 0$,

$$M^t_{1,v}(\psi) \geq e^{-2\varepsilon}\, M^t_{e^{-\varepsilon},u_i}(\psi_{-\varepsilon}) \frac{\bar{\mu}^f_{H^+}(B^+(u_i, e^{-\varepsilon}))}{\bar{\mu}^f_{H^+}(B^+(u_i, e^{\varepsilon}))}$$

Moreover, $M^t_{e^{-\varepsilon},u_i}(\psi_{-\varepsilon}) \to \int_{T^1M} \psi_{-\varepsilon} dm^f$ when $t \to +\infty$ (Theorem 7), whence:

$$\exists T > 0,\ \forall u_i,\ M^T_{e^{-\varepsilon},u_i}(\psi_{-\varepsilon}) \geq \frac{1}{2} \int_{T^1M} \psi_{-\varepsilon} dm^f.$$

We deduce that for all $v \in D$ (with $\varepsilon < \log 2$):

$$M^T_{1,v}(\psi) \geq \left(\frac{1}{16} \int_{T^1M} \psi \, dm^f \right) \inf_{w \in D} \frac{\bar{\mu}^f_{H^+}(B^+(w, \frac{1}{2}))}{\bar{\mu}^f_{H^+}(B^+(w,2))}$$

By Γ-invariance of the means, the above inequality is true for all $v \in \Lambda^2 \times \mathbf{R}$. Now, if $u \in \Lambda^2 \times \mathbf{R}$, we know that $M_{e^T,u}(\psi) = M^T_{1,\Phi^T u}(\psi)$ and above inequality applies with $v = \Phi^T u$. At last, we get

$$M^0_{e^T,u}(\psi)\, dM(u) \geq c \int_{T^1M} \psi dm^f, \quad \text{with} \quad c = \frac{1}{16} \inf_{w \in D} \frac{\bar{\mu}^f_{H^+}(B^+(w, \frac{1}{2}))}{\bar{\mu}^f_{H^+}(B^+(w,2))} > 0.$$

Finally, integrating with respect to M on D leads to (6) with $C = M(D)c$. This constant C does not depend on ψ, whence $m^f \ll M$. We deduce from the above that $\widehat{\mu}_o^f \ll \widehat{\nu}$. Indeed, let $A \subset \mathcal{H}$ be a Borel set, and assume that $\widehat{\nu}(A) = 0$. Let $0 < c < C$ be two constants. If A is small enough, we can choose for all $H \in A$ a distinguished vector $v \in H$, and a ray r chosen such that for all $H \in A$, $0 < c \leq \overline{\mu}_{H^+}^f(B^+(v,r) \leq C$. Then

$$\int_A d\widehat{\nu}(H) \int_H \mathbf{1}_{B^+(v,r)} \leq C\widehat{\nu}(A) = 0, \qquad \text{whence}$$
$$0 \leq c\widehat{\mu}_o^f(A) \leq \int_A d\widehat{\mu}_o^f(H) \int_H \mathbf{1}_{B^+(v,r)}\, d\overline{\mu}_{H^+v}^f = 0.$$

If A is not small enough, we can write it as a countable union of smaller sets. So we proved $\widehat{\mu}_o^f \ll \widehat{\nu}$.

Let us now prove that $\widehat{\mu}_o^f$ is ergodic: let $A \subset \mathcal{H}$ be a Borel set such that for all $\gamma \in \Gamma$, $\widehat{\mu}_o^f(\gamma A \triangle A) = 0$, and $\widehat{\mu}_o^f(A) > 0$. Then the measure $\widehat{\mu}_{o|A}^f$ is nonzero and quasi-invariant with cocycle $-c_o^f$. By what precedes, it satisfies $\widehat{\mu}_o^f \ll \widehat{\mu}_{o|A}^f$, and this implies $\widehat{\mu}_o^f(A^c) = 0$. So we have ergodicity.

Now let $\widehat{\nu}$ be a nonzero and quasi-invariant measure with cocycle c_o^f and let us show that it is also ergodic: if A is a Borel set in $\mathcal{H}$, Γ-invariant $\widehat{\nu}$-a.e, and if $\widehat{\nu}(A) > 0$, we can decompose the measure $\widehat{\nu}$ into $\widehat{\nu} = \widehat{\nu}_{|A} + \widehat{\nu}_{|A^c}$. By the above, there exist measurable maps ϕ and ϕ' such that $\widehat{\mu}_o^f = \phi\,\widehat{\nu} = \phi\,\widehat{\nu}_{|A} + \phi\,\widehat{\nu}_{|A^c}$ and $\widehat{\mu}_o^f = \phi'\,\widehat{\nu}_{|A}$. We deduce that $\widehat{\mu}_o^f(A^c) = 0$, and then necessarily that $\widehat{\nu}_{|A^c} = 0$, which means that $\widehat{\nu}$ is ergodic.

Moreover, the Radon-Nikodym derivative $\frac{d\widehat{\mu}_o^f}{d\widehat{\nu}}$ is Γ-invariant $\widehat{\nu}$-a.e, hence constant by ergodicity of $\widehat{\nu}$. So there exists a unique (up to a multiplicative constant) measure which is Γ-quasi-invariant with cocycle c_o^f.

Recall now that we consider the cohomology class on $\mathcal{H}$ of the Hölder cocycle c on $\partial\widetilde{M}$. Let $c' \in [c]_{\mathcal{H}} = [c_o^f]_{\mathcal{H}}$. There exists a positive Borel map φ on $\mathcal{H}$, such that for all $\gamma \in \Gamma$ and $(\xi, s) \in \mathcal{H}$, $c'(\gamma, (\xi, s)) = c_o^f(\gamma, \xi) + \varphi(\gamma(\xi, s)) - \varphi(\xi, s)$. If $\widehat{\nu}'$ is a Borel Radon measure on $\mathcal{H}$, which is Γ-quasi-invariant with the cocycle c' then an easy calculation shows that $\exp(-\varphi)\widehat{\nu}'$ is quasi-invariant with the cocycle c_o^f, it is then unique (up to a constant), and equals $\widehat{\mu}_o^f$, so $\widehat{\nu}'$ is also unique, and equals $\exp(\varphi)\widehat{\mu}_o^f$. $\square$

Proof of Theorem 2: It is clear that there exists a bijection between measures transverse to the strong unstable foliation on T^1M, which are quasi-invariant with the cocycle ρ^f, and measures transverse to the strong unstable foliation on $T^1\widetilde{M}$, which are Γ-invariant and quasi-invariant by holonomy with cocycle ρ^f. Let us consider the map: $M_T \to N_T$, which sends a Γ-invariant transverse measure, quasi-invariant by holonomy with cocycle ρ^f, to the measure $dN_T(v) = \exp(\rho_{v^-}^f(o, v))\, dM_T(v)$. It is clearly a bijection. An easy computation shows that N_T is a measure transverse to $\widetilde{\mathcal{F}}$ which is holonomy invariant and Γ-quasi-invariant with cocycle c_o^f. We can associate to N_T a Γ-quasi invariant measure

on $\mathcal{H}$ (with cocycle c_o^f), and this correspondance is 1-1. The existence of a unique quasi-invariant measure on $\mathcal{H}$ with cocycle c_o^f is then equivalent to the existence of a unique measure transverse to $\mathcal{F}$ with the cocycle ρ^f.
Theorem 2 then follows from Theorem 2 bis. $\square$

3.4 Equidistribution of means in a particular case

Lemma 3.3 means that the map $(u,r) \to \int_{B^+(u,r)} \psi \, d\bar{\mu}^f_{H^+}$ is "almost continuous". In this section, we shall see that under an additional assumption, it is really continuous, so that we can prove Theorem 3 directly, and deduce Theorem 2 bis as a corollary. The assumption is the following:

$$\forall u \in \Lambda^2 \times \mathbf{R}, \quad \bar{\mu}^f_{H^+}(\partial B^+(u,1)) = 0.$$

Note that it is true in the constant curvature case, and in the case of a surface, since the Patterson measure has no atoms.
With this assumption, we can prove that the means $(M^t_{1,u}(\psi))_{t\geq 0}$ are equicontinuous in $u \in \Lambda^2 \times \mathbf{R}$. We deduce that the convergence in Theorem 7 is uniform in $u \in \Lambda^2 \times \mathbf{R}$, and at last, relation (3) leads directly to the uniform convergence of the means $M_r(u)$ to $\int_{T^1M} \psi \, dm^f$ when $r \to \infty$.
Note that these two arguments of equidistribution of the means when $t \to +\infty$, and equicontinuity in $t \geq 0$ were already used in [12] in an algebraic framework.
Under the above assumption, Lemma 3.3 leads to:

Lemma 3.5 *If for all $u \in \Lambda^2 \times \mathbf{R}$, $\bar{\mu}^f_{H^+}(\partial B^+(u,1)) = 0$, then for all $u \in \Lambda^2 \times \mathbf{R}$, the family $(M^t_{1,u}(\psi))_{t\geq 0}$ is equicontinuous in $t \geq 0$.*

Sketch of the proof: Let $\varepsilon_0 > 0$, we shall find for all $u \in \Lambda^2 \times \mathbf{R}$ a neighbourhood $V = V(u, \varepsilon_0)$ of u, such that for all $v \in V$, for all $t \geq 0$, $|M^t_{1,v}(\psi) - M^t_{1,u}(\psi)| \leq \varepsilon_0$.
Since $\overline{\mu}^f_{H^+}(\partial B^+(u,1)) = 0$, there exists a $0 < \varepsilon \leq \varepsilon_0$, such that :

$$\left| \frac{\overline{\mu}^f_{H^+}(B^+(u, e^{\varepsilon}))}{\overline{\mu}^f_{H^+}(B^+(u, e^{-\varepsilon}))} \right| \leq e^{\varepsilon_0}. \tag{7}$$

Since ψ is continuous on Ω, one may choose ε small enough so that $\psi_\varepsilon \leq \psi + \varepsilon_0$. We put then $V(u, \varepsilon) = C(u, r_1, r_2, r_3)$: the cell given by Lemma 3.3 for ε. Now, relation (7) and Lemma 3.3, together with the fact that $[\psi \circ \Phi^t]_\varepsilon \leq \psi_\varepsilon \circ \Phi^t$ (since the geodesic flow expands unstable manifolds) allow to prove that for all $v \in V(u, \epsilon)$, and for all $t > 0$, $|M^t_{1,v}(\psi) - M^t_{1,u}(\psi)| \leq \varepsilon_0$. $\square$

With this Lemma, we can now prove the following equidistribution result for the horospherical means:

Theorem 3 bis: *Assume that for all $u \in T^1M$ $\bar{\mu}^f_{H^+}(\partial B^+(u,1)) = 0$, then for all continuous map $\psi : T^1M \to \mathbf{R}$, the means $M_{r,u}(\psi)$ converge uniformly in $u \in \Lambda^2 \times \mathbf{R}$ to $\int_{T^1M} \psi \, dm^f$.*

Proof: We consider now the means as functions on T^1M. By the above Lemma, for all $u \in \Lambda^2 \times \mathbf{R}$, the means $(M^t_{1,u})_{t\geq 0}$ are equicontinuous, whence (Theorem 7) it is easy to prove that they converge uniformly on Ω to $\int_{T^1M} \psi\, dm^f$ when $t \to +\infty$. Let $\varepsilon > 0$, there exists a $T > 0$, such that:

$$\forall\, t \geq T,\ \forall\, u \in \Omega, \qquad \left| M^t_{r,u}(\psi) - \int_{T^1M} \psi\, dm^f \right| \leq \varepsilon.$$

Let now $v \in \Omega$ be a vector, and $t \geq T$; as Ω is invariant by the flow, the above inequality applies for $u = \Phi^{-t}v$ and above relation (3) gives the desired result:

$$\left| M^0_{re^t,v}(\psi) - \int_{T^1M} \psi\, dm^f \right| \leq \varepsilon.$$

$\square$

We can now easily deduce Theorem 2 bis from this result:

Sketch of the proof: We use the equality of Lemma 3.4. By the uniform equidistribution property of means, the left term converges to $M(D)\int_{T^1M} \psi\, dm^f$, and the right term is smaller than $N \int_{T^1M} \psi\, dM$. Then we have directly $m^f \ll M$, and we conclude the proof as in the general case. $\square$

4 Growth of leaves of the horospherical foliation, equidistribution of horospheres

The aim of this section is to prove Theorem 3. In fact, using Theorem 2 bis, we shall prove a more general result of equidistribution of sets on leaves satisfying a certain condition of growth, denoted by $(*)$ (Theorem 8). We deduce Theorem 3 as a corollary, since the horospherical balls have a polynomial growth (Lemma 4.2), whence they satisfy $(*)$ (Lemma 4.1).

This paragraph is inspired by Plante [25], and Goodman-Plante [14]. These two references define a notion of "averaging sets" for the foliation, i.e. sequences of sets on transversals to the foliation, defined so that they become equidistribued to a holonomy invariant measure. The idea here is to introduce an analogous notion to obtain equidistribution to the holonomy quasi-invariant measure $\{\mu^f_T\}$. But it is more appropriate to work with the holonomy invariant measure $\{\bar{\mu}^f_T\}$ on transversals to $\widetilde{\mathcal{W}}^{su}$. The definition of an averaging set in [14] involves the holonomy pseudogroup of the foliation. In our case, this pseudogroup is trivial on $\widetilde{\mathcal{W}}^{su}$. Moreover, the above transverse measure is Γ-quasi-invariant, so we will introduce a condition dealing with Γ.
Let us now fix a sequence $(E_n)_{n\in\mathbf{N}}$ of compact sets of a horosphere H^+. If they satisfy the condition $(*)$ described below, then we shall prove (Theorem 8) that the means on the E_n w.r.t. the measure $\bar{\mu}^f_{H^+}$ become equidistribued to m^f.

Let D be any proper fundamental domain for the action of Γ on $T^1\widetilde{M}$, that is an open set D, such that $D = \overset{\circ}{\bar{D}}$, $\cup_{\gamma\in\Gamma}\gamma\bar{D} = T^1\widetilde{M}$, and the $(\gamma D)_{\gamma\in\Gamma}$ are pairwise disjoint. For fixed $n \in \mathbf{N}$, we distinguish three types of elements $\gamma \in \Gamma$:

1- First, the set $\Gamma_n^1(D)$ of elements γ such that E_n “crosses completely” γD, i.e.

$$E_n \cap \gamma D = H^+(u) \cap \gamma D;$$

2- Second, the set $\Gamma_n^2(D)$ of elements γ such that E_n crosses partly γD, or again

$$\emptyset \neq E_n \cap \gamma D \subsetneq H^+(u) \cap \gamma D$$

3- Third, the set $\Gamma_n^3(D)$ of elements γ such that $\gamma D \cap E_n = \emptyset$.

Compacity of the E_n implies finiteness of the sets $\Gamma_n^1(D)$ and $\Gamma_n^2(D)$.
Condition $(*)$ is then the following: there exists a fundamental domain D, such that, with the above notations:

$$\lim_{n\to\infty} \frac{1}{\bar{\mu}^f_{H^+}(E_n)} \sum_{\gamma\in\Gamma_n^2(D)} \bar{\mu}^f_{H^+}(E_n \cap \gamma D) \;=\; 0 \qquad (*)$$

In other words, we can forget the "boundary terms" $\Gamma_n^2.D$: only the images γD of D completely crossed by E_n must be taken into account.
We can now state the following:

Theorem 8 *Let $(E_n)_{n\in\mathbf{N}}$ be a sequence of compact sets on the horosphere $H^+(u)$, with $u \in \Lambda^2 \times \mathbf{R}$ and $\cup E_n = H^+(u)$. If the sequence $(E_n)_{n\in\mathbf{N}}$ satisfy $(*)$, then the sequence*

$$dM_n(w) \;=\; \frac{1}{\bar{\mu}^f_{H^+}(E_n)}\, \mathbf{1}_{E_n}(w)\, d\bar{\mu}^f_{H^+}(w)$$

converges weakly to m^f when $n \to +\infty$.

We will prove this result at the end of this section. Let us first apply it to sequences of balls $B^+(u,r)$, when $r \to \infty$. We shall first prove that balls satisfy $(*)$, and thus get the equidistribution result stated in the introduction (Theorem 3) as an immediate corollary of Theorem 8.

Lemma 4.1 *The sequence of balls $B^+(u,r)$ satisfies $(*)$.*

As a straightforward corollary, Theorem 3 is proved by the two above results.

Proof: Let D be any fundamental domain. By compacity of $\bar{D}$, we can find a constant $r_0 > 0$, such that for all $v \in D$, $B^+(v, r_0) \cap D = H^+(v) \cap D$. Now if $\emptyset \neq B^+(u,r) \cap \gamma D \subsetneq H^+(u) \cap \gamma D$, then triangular inequality gives $B^+(u, r + r_0) \cap \gamma D = H^+(u) \cap \gamma D$. Moreover, the same argument gives also

$B^+(u, r - r_0) \cap \gamma D = \emptyset$. This proves that $B^+(u,r) \cap \Gamma_r^2 D \subset B^+(u, r+r_0) \setminus B^+(u, r - r_0)$. By Lemma 4.2, it is easy to prove that

$$\lim_{r \to +\infty} \frac{1}{\bar{\mu}^f_{H^+}(B^+(u,r))} \sum_{\gamma \in \Gamma_r^2} \bar{\mu}^f_{H^+}(B^+(u,r) \cap \gamma D) = 0.$$

$\square$

Lemma 4.2 *For all fixed $u \in T^1\widetilde{M}$, $\bar{\mu}^f_{H^+(u)}(B^+(u,r)) = O(r^{2\delta^f + 4\|f\|_\infty})$.*

Proof: If we consider $\Phi^t(u)$ instead of u, we can assume that the origin o is in the interior of the horoball defined by u^- and u, or again $\beta_{u^-}(o,u) \leq 0$. As $\nu_o^f(\partial\widetilde{M}) = 1$, and by definition of the measure $\bar{\mu}^f_{H^+(u)}$, we have

$$\bar{\mu}^f_{H^+(u)}(B^+(u,r)) \leq \sup_{v \in B^+(u,r)} \left(\exp(\delta^f \beta_{v^+}(o,v) - \rho^f_{v^+}(o,v) - \rho^f_{v^-}(o,v)) \right).$$

Let us first find an upper bound for the quantity $|\rho^f_{v^+}(o,v) + \rho^f_{v^-}(o,v)|$. Since the curvature of $\widetilde{M}$ is negative, we can find a triangle (a,b,c) such that $a \in [o, v^+)$, $b \in [o, v^-)$, $c \in (v^-, v^+)$, which satisfies: $d(o,a) = d(o,b)$, $\beta_{v^+}(a,c) = 0$ and $\beta_{v^-}(b,c) = 0$, and also $d(a,b) \leq \delta$, $d(a,c) \leq \delta$, $d(b,c) \leq \delta$, with δ a constant which depends only on the upper bound of the sectional curvature of M. Now, as f is Hölder, with exponent $\alpha(f)$, we can easily check that:

$$|\rho^f_{v^+}(o,v) + \rho^f_{v^-}(o,v)| \leq 2\|f\|_\infty \, d(o,a) + C(f,\delta).$$

Moreover, $d(o,a) \simeq d(o,c) \simeq d(o, (v^-, v]) \leq d(o,v)$, as o is in the interior of the horoball defined by v and v^-. For the same reason, and by convexity of horoballs, the angle at v between $[o,v]$ and $[v, v^+)$ is greater than $\pi/2$, whence $d(o,v) \simeq \beta_{v^+}(o,v)$. We deduce

$$\bar{\mu}^f_{H^+(u)}(B^+(u,r)) \leq Cst. \sup_{B^+(u,r)} \exp((\delta^f + 2\|f\|_\infty).\beta_{v^+}(o,v)).$$

As the Busemann cocycle is continuous, there exists $\alpha > 0$, such that $d_o(u^+, v^+) \leq \alpha$ implies $|\beta_{v^+}(o,v)| \leq \beta_{u^+}(o,u) + 1$. Moreover, by definition of d_{H^+}, if $d_o(u^+, v^+) \geq \alpha$, we have the following bound:

$$\exp(\frac{1}{2}\beta_{v^+}(o,v)) \leq \frac{1}{\alpha} \, d_{H^+}(u,v) \, \exp(-\beta_{u^+}(o,u)/2) \leq C(u).r,$$

and at last, if $v \in B^+(u,r)$, we have:

$$e^{(\delta^f + 2\|f\|_\infty)\beta_{v^+}(o,v)} \leq \max\left(e^{(\delta^f + 2\|f\|_\infty).(\beta_{u^+}(o,u)+1)}, \; C(u).r^{2\delta^f + 4\|f\|_\infty} \right).$$

All these inequalities give the desired comparison (if u is fixed). $\square$

Let us now prove Theorem 8:

Proof of Theorem 8: It is enough to show convergence of means for small Borel sets $B \subset D \subset T^1M$ which can be written on the form $B = T \times b$ in a chart of the foliation, with T a transversal to $\mathcal{W}^{su}$, and b a "piece" of leaf. Lift such a Borel set B into $\widetilde{B}_0 = \widetilde{T}_0 \times \widetilde{b}_0 \subset D$, and set $\widetilde{B} = \cup_{\gamma\in\Gamma}\gamma\widetilde{B}_0$. We can write again the means on the following form:

$$\begin{aligned}
M_n(B) &= \frac{1}{\bar{\mu}^f_{H^+}(E_n)} \sum_{\gamma\in\Gamma} \int_{H^+(u)} \mathbf{1}_{E_n\cap\gamma\widetilde{B}_0}(v)\, d\bar{\mu}^f_{H^+}(v) \\
&= \int_{\widetilde{T}_0} \frac{1}{\bar{\mu}^f_{H^+}(E_n)} \sum_{\gamma\in\Gamma} e^{\rho^f_{v^-}(o,\gamma^{-1}o)} \mathbf{1}_{\widetilde{T}_0\cap\gamma^{-1}E_n\neq\emptyset}(v) \bar{\mu}^f_{H^+}(H^+(v)\cap\widetilde{B}_0) \\
&+ R(n,\widetilde{T}_0), \\
&= \int_{\widetilde{T}_0} d\nu^n_{\widetilde{T}_0}(v) \int_{H^+(v)} \mathbf{1}_{\widetilde{B}_0}(w)\, d\bar{\mu}^f_{H^+}(w) + R(n,\widetilde{T}_0) \qquad \text{with} \\
d\nu^n_{\widetilde{T}_0}(w) &= \frac{1}{\bar{\mu}^f_{H^+}(E_n)} \sum_{\gamma\in\Gamma} \exp(\rho^f_{v^-}(o,\gamma^{-1}o)) \mathbf{1}_{\widetilde{T}_0\cap\gamma^{-1}E_n\neq\emptyset}(w),
\end{aligned}$$

and the rest $R(n,\widetilde{T}_0)$ corresponds to the $\gamma \in \Gamma$ such that E_n crosses only partly the box $\gamma\widetilde{B}_0$; so it is bounded by:

$$R(n,\widetilde{T}_0) \leq \frac{1}{\bar{\mu}^f_{H^+}(E_n)} \sum_{\gamma\in\Gamma^2_n} \bar{\mu}^f_{H^+}(E_n \cap \gamma D).$$

The sequence (M_n) of probabilities is defined on the compact set Ω, so it has limit points for the weak topology, and after the above computation, for any fixed transversal T, these limit points are in correspondance $1-1$ with those of the sequence (ν^n_T).

Let M be a limit point of measures M_n; if we consider a subsequence, we can assume that the M_n converge to M, and the measures ν^n_T also converge to a measure ν_T. This family of measures is invariant by holonomy; indeed, if we consider two small transversals T and T' to the foliation, such that $B = T \times b = T' \times b$ in a chart of the foliation, then

$$\begin{aligned}
M(B) &= \lim_{n\to\infty} \int_T d\nu^n_T(v) \int_{H^+(v)} \mathbf{1}_{\widetilde{B}_0}(w) d\bar{\mu}^f_{H^+}(w) \\
&= \lim_{n\to\infty} \int_{T'} d\nu^n_{T'}(v) \int_{H^+(v)} \mathbf{1}_{\widetilde{B}_0}(w) d\bar{\mu}^f_{H^+}(w),
\end{aligned}$$

which gives invariance by holonomy of the limit measures. This family of measures is Γ-quasi-invariant; indeed, if $g \in \Gamma$, it is easy to compute that:

$$\begin{aligned}
d\nu^n_{gT}(gv) &= \frac{1}{\bar{\mu}^f_{H^+}(E_n)} \sum_{\gamma\in\Gamma} \exp(\rho^f_{gv^-}(o,\gamma^{-1}o))\, \mathbf{1}_{gT\cap\gamma^{-1}E_n}(gv) \\
&= \exp(\rho^f_{v^-}(g^{-1}o,o))\, d\nu^n_T(v) \;=\; \exp(c^f_o(g,v^-))\, d\nu^n_T(v).
\end{aligned}$$

At last, we proved that a limit point M of M_n induces a transverse measure invariant by holonomy and quasi-invariant by Γ with cocycle c_o^f, whence (Theorem 2) it equals the transverse measure $(\bar{\mu}_T^f)_T$ induced by m^f; this shows that the sequence M_n becomes equidistributed to m^f. $\square$

5 Application to coverings

5.1 Nonsymmetric potential

If f is not symmetric, we set $\check{f}(v) = f(-v)$, and we look again at the product measure $\nu_o^f \times \nu_o^{\check{f}}$ on $\partial^2 \widetilde{M}$. We are still in the case when $\delta^f < +\infty$, and we have $\delta^f = \delta^{\check{f}}$ (see [9]). The equilibrium state m^f is still constructed by means of this product, but the density is slightly different [9]:

$$d\widetilde{m}^f(v^-, v^+, s) = ds\, d\nu_o^f(v^+)\, d\nu_o^{\check{f}}(v^-)\ \exp\left(\beta_{v^+}^f(o,v) + \beta_{v^-}^{\check{f}}(o,v)\right).$$

The measures of paragraph 2.2 can then be written:

$$d\widehat{\mu}_o^f(\xi, s) = ds\, d\nu_o^{\check{f}}(v^-)\ \exp(-\delta^f s), \qquad \text{and}$$

$$d\overline{\mu}_{H^+u}^f(v) = d\nu_o^f(v^+)\ \exp(\beta_{v^+}^f(o,v) - \rho_{v^-}^{\check{f}}(o,v)).$$

And the relations of Γ-quasi-invariance become:

$$\frac{d(\gamma_*^{-1}\widehat{\mu}_o^f)}{d\widehat{\mu}_o^f}(\xi, s) = \exp(c_o^{\check{f}}(\gamma, \xi)), \quad \text{and} \quad \frac{d(\gamma_*^{-1}\overline{\mu}_{H^+(\gamma v)}^f)}{d\widehat{\mu}_{H^+v}^f}(v) = \exp(-\rho_{v^-}^{\check{f}}(\gamma^{-1}o, o))$$

All proofs apply in the same way, since they are based on the Hölder continuity of f and geometry of M, whence Theorems 2 and 2 bis are still true.

5.2 Proof of Theorem 4

The above paragraph applies in particular to the case of a closed 1-form α on M. Indeed, a 1-form on M is a smooth map from M into T^*M which to each $x \in M$ associates a linear form α_x on T_xM. In particular, it is a map from TM into $\mathbf{R}$, that we can restrict to T^1M. For all $x \in M$, linearity of α_x gives antisymmetry of α: $\alpha(-v) = -\alpha(v)$. Moreover, as α is closed, we have $\rho_\xi^{\check{\alpha}}(x,y) = \int_x^y \check{\alpha} = -\rho_\xi^\alpha(x,y)$, whence (cf [3]):

$$d\widetilde{m}^\alpha(v^-, v^+, s) = ds\, d\nu_o^\alpha(v^+)\, d\nu_o^{\check{\alpha}}(v^-)\ \exp\left(\delta^\alpha \beta_{v^-}(o,v) + \delta^\alpha \beta_{v^+}(o,v)\right).$$

The measure $\widehat{\mu}_o^\alpha$ can be written: $d\widehat{\mu}_o^\alpha(u^-, s) = ds\, d\nu_o^{\check{\alpha}}(v^-)\ \exp(-\delta^\alpha s)$. Moreover, since α is closed, we have $c_o^{\check{\alpha}}(\gamma, \xi) = \rho_\xi^{\check{\alpha}}(\gamma^{-1}o, o) = -\int_o^{\gamma o} \alpha$, whence:

$$\frac{d(\gamma_*^{-1}\widehat{\mu}_o^\alpha)}{d\widehat{\mu}_o^\alpha}(\xi, s) = \exp(-\int_o^{\gamma o} \alpha). \tag{8}$$

Remark that $\gamma_*^{-1}\widehat{\mu}_o^\alpha$ is a multiple of $\widehat{\mu}_o^\alpha$ and does not depend on o: the cocycle does not depend on ξ and o, but only on γ, and the measure ν_o^α on $\partial\widetilde{M}$ is independent on o too. So we will omit the subscript o for these measures $\widehat{\mu}^\alpha$ et ν^α.

Let us recall some notions: the first singular homology group $H_1(M, \mathbf{Z})$ over $\mathbf{Z}$ can be identified to the abelianization $\Gamma/[\Gamma,\Gamma]$ of the fundamental group. The first real homology group $H_1(M,\mathbf{R})$ is the $\mathbf{R}$-vector space generated by $H_1(M,\mathbf{Z})$ except the torsion elements: $H_1(M,\mathbf{R}) = H_1(M,\mathbf{Z}) \otimes_{\mathbf{Z}} \mathbf{R}$. The first de Rham cohomology group $H^1_{dR}(M)$ is the set of closed 1-forms modulo exact forms. Moreover, the de Rham isomorphism (see [28]) allows to identify the dual of $H_1(M,\mathbf{R})$ with $H^1_{dR}(M)$. This identification is induced by the map:

$$\begin{array}{rcl} H_1(M,\mathbf{R}) \times H^1_{dR}(M) & \to & \mathbf{R} \\ (\gamma,\alpha) & \mapsto & \displaystyle\int_\gamma \alpha\,. \end{array}$$

We denote by $\int_\gamma \alpha$ the integral of any representant of α on any loop of M belonging to the homology class of γ. In particular, it is the integral of α on the unique closed geodesic in the homotopy class of γ, and it is also the integral $\int_o^{\gamma o} \alpha$, since by definition of the action of Γ on $\widetilde{M}$, the curve $[o,\gamma o]$ projects on a loop in the class γ.
Theorem 2 bis allows to determine the measures on $\mathcal{H}$, which are invariant by a normal subgroup $\overline{\Gamma}$ of Γ, and quasi-invariant by Γ. In other words, if $\overline{M}$ is a regular cover of M, with fundamental group $\overline{\Gamma}$, we can determine the invariant measures on the space of horospheres of $\overline{M}$, which are quasi-invariant by the group $\Gamma/\overline{\Gamma}$ of the cover. If α is a closed 1-form which vanishes on the image of $\overline{\Gamma}$ in $H_1(M,\mathbf{R})$, formula (8) above shows that the measure $\widehat{\mu}^\alpha$ induced by the Gibbs measure m^α is in particular invariant by $\overline{\Gamma}$. Theorem 4 establishes the converse statement under the additional assumption: if $\widehat{\nu}$ is $\overline{\Gamma}$-invariant, $\overline{\Gamma}$-ergodic, and quasi-invariant by Γ, then it is necessarily of the form $\widehat{\nu} = \widehat{\mu}^\alpha$, with α a 1-form on M vanishing on loops of $\bar{\Gamma}$.

Proof of Theorem 4: The first part of the Proposition was proved above. The group $\overline{\Gamma}$ is normal in Γ, so the action of an element $a \in A = \Gamma/\overline{\Gamma}$ on $\widehat{\nu}$ makes sense: indeed, we can write $a = \gamma_a\overline{\Gamma} = \overline{\Gamma}\gamma_a$, and $\overline{\Gamma}$-invariance of $\widehat{\nu}$ shows that if $\gamma_a \in \Gamma$, the measure ${\gamma_a^{-1}}_*\widehat{\nu}$ only depends on $a \in A$. It shows also that for all $a \in A$, the measure $a_*^{-1}\widehat{\nu}$ is also $\overline{\Gamma}$-invariant. The derivative $\frac{da_*^{-1}\widehat{\nu}}{d\widehat{\nu}}$ is then a $\overline{\Gamma}$-invariant map $\widehat{\nu}$-a.e, and by ergodicity of $\widehat{\nu}$, it is constant. So for all $a \in A$, there exists a constant $\lambda(a) > 0$, such that $a_*^{-1}\widehat{\nu} = \lambda(a)\widehat{\nu}$. The map λ is clearly a morphism of groups from A into $\mathbf{R}_+^*$, and when composed with the quotient morphism $\Gamma \to A$, it defines a homomorphism, still denoted by λ, from Γ to $\mathbf{R}_+^*$. Moreover, as $\mathbf{R}_+^*$ is abelian, λ induces a morphism defined on the abelianization $\Gamma/[\Gamma,\Gamma] = H_1(M,\mathbf{Z})$. We can then extend it by $\mathbf{R}$-linearity into the exponential of a linear form on $H_1(M,\mathbf{R})$. The de Rham isomorphism between $H_1(M,\mathbf{R})$ and $H^1_{dR}(M)$ allows to see $\log\lambda$ as an element of $H^1_{dR}(M)$; so let $-\alpha$ be a closed 1-form in its class; for all loop $c(\gamma)$ in the homology class of γ,

$\lambda(\gamma) = \exp(-\int_{c(\gamma)} \alpha)$. In particular, $\lambda(\gamma) = \exp(-\int_{\pi[o,\gamma o]} \alpha) = \exp(-\int_o^{\gamma o} \alpha)$. Moreover, a closed 1-form α is in particular a Hölder map on the unit tangent bundle T^1M, hence we have: for all $\gamma \in \Gamma$, $\gamma_*^{-1}\widehat{\nu} = \exp(-\int_o^{\gamma o} \alpha)\,\widehat{\nu}$. Then Theorem 2 bis applies, and $\widehat{\nu}$ is necessarily equal to $\widehat{\mu}^{\alpha}$. $\square$

References

[1] J. Aaronson, *An introduction to infinite ergodic theory* (1997) American Math.Soc. Mathematical Surveys and Monographs, Vol 50.

[2] M. Babillot, *On the mixing property for hyperbolic systems* (2001) To appear in Israël J. of Maths.

[3] M. Babillot, *Géodésiques et horocycles sur le revêtement d'homologie d'une surface hyperbolique*, Séminaire théorie spectrale et géométrie, Grenoble, (1996), 89-104

[4] M. Babillot, F. Ledrappier, *Geodesic paths and Horocycle Flow on Abelian Covers* (1998) Proc. International Colloquium on Lie groups and Ergodic theory, Tata Institute of Fundamental Research, Narosa Publishing house, New Delhi, 1-32.

[5] M. Bourdon, *Structure conforme au bord et flot géodésique d'un CAT(-1)-espace* (1995), L'Enseignement Mathématique, 41, 63-102.

[6] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture Notes in Mathematics, Vol. 470. Springer-Verlag, Berlin-New York (1975).

[7] R. Bowen, B. Marcus, *Unique ergodicity for horocycle foliations* (1977), Israel J. of Maths. **26**, 43-67.

[8] A. Connes, *Sur la théorie non commutative de l'intégration*, in Algèbres d'Opérateurs, Séminaire, Les Plans-sur-Bex, Suisse, (1978) Lecture Notes in Mathematics, Vol. 725, Springer-Verlag, Berlin Heidelberg New York.

[9] Y. Coudène, *Gibbs measures on negatively curved manifolds* (2001) Prépublication de l'E.N.S.

[10] F. Dal'bo, *Topologie du feuilletage fortement stable* (2000), Ann. Inst. Fourier, 50, N.3, 981-993.

[11] P. Eberlein, *Geometry of nonpositively curved manifolds* (1996), Chicago Lectures in Mathematics.

[12] R. Ellis, W. Perrizo, *Unique ergodicity of flows on homogeneous spaces*, (1978), Israël Journal of Maths, Vol 29, N 2-3, 276-284.

[13] H. Furstenberg, *The unique ergodicity of the horocycle flow*, Springer Lecture Notes in Math, **318** (1972), 95-115.

[14] S.E. Goodman, J.F. Plante, *Holonomy and averaging in foliated sets*, J.Differential Geometry, **14** (1979), n.3, 401-407.

[15] U. Hamenstädt, *A new description of the Bowen-margulis measure* (1989), Ergodic Theory and Dynamical Systems, 9, 455-464.

[16] U. Hamenstädt, *Cocycles, Hausdorff measures and cross ratios* (1997), Ergodic Theory and Dynamical Systems, 17, 1061-1081.

[17] V. Kaimanovitch, *Ergodicity of harmonic invariant measures for the geodesic flow on hyperbolic spaces*, J.Reine Angew.Math 455 (1994), 57-103.

[18] V. Kaimanovitch, *Invariant measures of the geodesic flow and measures at infinity on negatively curved manifolds* (1990), vol 53, n.4, 361-393.

[19] V. Kaimanovich, *Ergodic properties of the horocycle flow and classification of Fuchsian groups*, J. Dynam. Control Systems 6 (2000), no. 1, 21–56.

[20] V. Kaimanovich, *SAT Actions and Ergodic Properties of the Horosphere Foliation* (2001), preprint.

[21] G. Keller, *Equilibrium States in Ergodic theory*, London Math. Soc. Student Texts, **42**, Cambridge University Press, Cambridge (1998).

[22] F. Ledrappier, *Structure au bord des variétés à courbure négative*, Séminaire de théorie spectrale et géométrie, Grenoble (1994-95), 93-118.

[23] J-P. Otal, *Sur la géométrie symplectique de l'espace des géodésiques d'une variété à courbure négative*, Rev. Math. Iber. **8** (1992) 441-456.

[24] S.J. Patterson, *The limit set of a Fuchsian Group*, Acta Math. **136** (1976), 241-273.

[25] J.F. Plante, *Foliations with measure preserving holonomy*, Annals of Maths, **102** (1975), 327-361.

[26] T. Roblin, *Ergodicité et unique ergodicité du feuilletage horosphérique, mélange du flot géodésique et équidistributions diverses dans les groupes discrets en courbure négative* (2001), Preprint.

[27] P. Walters, *An Introduction to ergodic theory*, G.T.M. 79, Springer Berlin (1982)

[28] F.W. Warner, *Foundations of Differentiable manifolds and Lie Groups* (1971), Scott, Foresman and Company, Glenview, Illinois.